

Soil warming changes litter chemistry and fungal community composition but not decomposition rate

Eric W. Morrison,^{1*} Linda T. A. van Diepen,¹ Serita D. Frey,¹ Anne Pringle,² A. Stuart Grandy,¹ Chris M. Sthultz³

1. Department of Natural Resources and the Environment, University of New Hampshire; 2. Department of Botany, University of Wisconsin; 3. Math, Science and Technology Department, University of Minnesota



Background

Increased temperatures are expected to increase rates of litter decay, but temperature effects on fungal communities will likely mediate this response. We used an *in situ* litter decay experiment to examine effects of warming on fungal community composition, enzyme activities, and litter chemistry in decomposing litter.

Methods

• Leaf litter bags were allowed to decompose for one or two years in a forest stand exposed to 9 years of soil warming (+5 °C above ambient).

• RNA and DNA were extracted from litter, and the fungal ITS2 region was sequenced by 454 sequencing.

• Extracellular enzyme activities were measured using standard methods:

- Beta-glucosidase (BG)
- Cellobiohydrolase (CBH)
- Phenol oxidase (PPO, ABTS)
- Peroxidase (PERgross, TMB)
- Leucine aminopeptidase (LAP)
- N-acetylglucosaminidase (NAG)
- Acid-phosphatase (PHOS)

• Molecular litter chemistry was measured by py-GC/MS.

Treatment abbreviations


- Control: ambient temperature
- 5C: heated 5 °C above ambient




*Presenting author email: eric.morrison@unh.edu

Results


Warming did not affect decay rate, but did reduce litter quality




Altered enzyme activities after two years coincided with altered fungal community composition




ECM fungi were negatively correlated with saprotrophs and BG:PPO



Russulaceae were favored by warming



ECM fungi were favored by warming



Summary

Warming increased lignin relative abundance and decreased BG:PPO after two years.

A shift in enzyme activities was correlated with a concomitant shift in the fungal community.

Warming has increased tree growth and N mineralization at this site (Melillo *et al.* 2011, *PNAS*). ECM fungi may be favored by increased photosynthate exudation.

Competition between ECM fungi and saprotrophs may limit decay rate.

Acknowledgements

This work was supported by the NSF Long-term Ecological Research (LTER) Program and an NSF Ecosystem Studies grant (DEB-1021063) to SDF and AP. We would especially like to acknowledge the NSF LTER program for providing intellectual and financial support of long-term ecological experiments without which this work would not have been possible. Sequencing was performed at the University of Illinois Roy J. Carver Biotechnology Center.