



Bursty Bulk Flows: Comparing THEMIS Observations with OpenGGCM Simulations.

Derick M. Arel, Joachim Raeder, Bashi Ferdousi, Kristofor Maynard, W. Douglas Cramer
 University of New Hampshire, Space Science Center, Durham NH


Themis Observations 2009

THEMIS observations of Bursty Bulk Flow (BBF) events in the Earth's magnetotail feature varying magnetic field (B_z) signatures for the same BBF, such as a dip in B_z which precedes the flow, as it propagates earthward through the central plasma sheet. It has not been clear whether these variations are due to some aspect of the BBF that is changing in time or due to the geometry of the BBF and the specific location of the spacecraft relative to it.




Figures from: Runov, A., V. Angelopoulos, M. I. Sitnov, V. A. Sergeev, J. Bonnell, J. P. McFadden, D. Larson, K.-H. Glassmeier, and U. Auster (2009). THEMIS observations of an earthward-propagating dipolarization front. *Geophys. Res. Lett.*, 36, L14106, doi:10.1029/2009GL038980.

OpenGGCM Simulation with virtual satellites.



Virtual observations of the simulated BBF.



In order to investigate these varying BBF signatures, we use OpenGGCM simulations, which produce realistic looking BBFs. By inserting an array of virtual satellites, we generate virtual observations at different locations relative to the BBFs. We show that the same BBF produces highly varying signatures depending on the location of the observing satellite. Specifically we show that the dip in B_z seems to depend on where the satellite is relative to the BBF. This results mainly from the fact that BBFs are not simple one-dimensional structures as they are often depicted in the literature, but rather following more complicated paths in the plasma sheet.

Left: The small circles represent the locations of virtual satellites.

Left: Data collected by virtual satellites within the simulation.

Above: A zoomed-in image of the B_z data collected as the simulated BBF passed the virtual satellite array. The negative B_z dip is a feature found on the $-Y$ side of the BBF, but it is not prominent, or is non-existent, on the $+Y$ side of the BBF. This demonstrates a variation in B_x signatures similar to those observed by the THEMIS spacecraft.

Conclusion: The observed signature depends on a spacecraft's position relative to the BBF.